

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 39/00		A2	(11) International Publication Number: WO 99/33488
			(43) International Publication Date: 8 July 1999 (08.07.99)
(21) International Application Number: PCT/EP98/08562		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 18 December 1998 (18.12.98)		Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(30) Priority Data: 9727262.9 24 December 1997 (24.12.97) GB			
(71) Applicant (for all designated States except US): SMITHK-LINE BEECHAM BIOLOGICALS S.A. [BE/BE]; Rue de l'Institut 89, B-1330 Rixensart (BE).			
(72) Inventors; and (75) Inventors/Applicants (for US only): DALEMANS, Wilfried, L., J. [BE/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE). LAFERRIERE, Craig, Antony, Joseph [CA/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE). PRIEELS, Jean-Paul [BE/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE).			
(74) Agent: DALTON, Marcus, Jonathan, William; SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).			
(54) Title: VACCINE			
(57) Abstract The present invention provides vaccine formulations comprising T-independent or polysaccharide conjugate vaccines adjuvanted with an immunostimulatory CpG oligonucleotide.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Vaccine

The present invention relates to new vaccine formulations, and to methods
5 for their production and their use in medicine.

Immunomodulatory oligonucleotides containing unmethylated CpG
dinucleotides ("CpG") are known (WO 96/02555, EP 468520). CpG is an
abbreviation for cytosine-guanosine dinucleotide motifs present in DNA.
Historically, it was observed that the DNA fraction of BCG could exert an anti-
10 tumor effect. In further studies, synthetic oligonucleotides derived from BCG gene
sequences were shown to be capable of inducing immunostimulatory effects (both in
vitro and in vivo). The authors of these studies concluded that certain palindromic
sequences, including a central CG motif, carried this activity (Tokunaga, T. et al.
Microbial. Immunol. 36: 55 (1992)). The central role of the CG motif in
15 immunostimulation was later elucidated in a publication by Krieg (Nature 374 p546
1995). Detailed analysis has shown that the CG motif has sequences that are
common in bacterial DNA but are rare in vertebrate DNA.

It is currently believed that this evolutionary difference allows the vertebrate
immune system to detect the presence of bacterial DNA (as occurring during an
20 infection) leading consequently to the stimulation of the immune system.
Immunostimulatory activity has been shown for sequences as small as 15 nucleotide
bases (Krieg, et al. Nature 374: 546 (1995)) and that the CpG motif has to be
unmethylated. It has been postulated that the oligo should be in a hexamer setting:
purine purine CG pyrimidine pyrimidine, but this is not obligatory.

25 Streptococcus pneumoniae is a gram positive bacteria that is pathogenic for
humans, causing invasive diseases such as pneumonia, bacteremia and meningitis,
and diseases associated with colonisation, such as acute Otitis media. The
mechanisms by which pneumococci spread to the lung, the cerebrospinal fluid and
the blood is poorly understood. Growth of bacteria reaching normal lung alveoli is
30 inhibited by their relative dryness and by the phagocytic activity of alveolar
macrophages. Any anatomic or physiologic derangement of these co-ordinated
defenses tends to augment the susceptibility of the lungs to infection. The cell-wall
of Streptococcus pneumoniae has an important role in generating an inflammatory

response in the a veoli of the lung (Gillespie et al , I&I 65: 3936). The release of cell-wall components occurs at the end of the pneumococcal growth cycle by autolysis due to the synthesis of the protein N-acetyl muramoyl-L-alanine amidase (lytA). DNA will also be released into the infected region upon autolysis of the pneumococci.

In order for the organism to have an effective immune response against invading bacteria, it must have mechanisms to coordinate the type of immune response most likely to stop infection. For intracellular pathogens, the coordination appears to occur between cell mediated or humoral immune responses, and these are controlled by T-cells of the type Th1 and Th2. However, extracellular bacteria frequently employ a polysaccharide either in the form of a capsule or a lipopolysaccharide to protect themselves from the effects of serum complement which can lyse the bacteria, or render them accessible to phagocytes such as macrophage and neutrophils.

In this case, the immune response follows another path, the T-independent immune response. The T-independent immune response may be further divided into Type 1 and Type 2. T-independent type 2 antigens possess the characteristics embodied by polysaccharide antigens, including: large molecular weight, repeat antigenic epitopes, ability to activate the complement cascade, poor in vivo degradability and inability to stimulate MHC class II dependent T cell help (Mond et al. Annu Rev Immunol 13:655-92). The Type 1 antigens, unlike the polysaccharides, are mitogenic for B-cells, and are comprised of the lipopolysaccharides (LPS). T-independent Type 2 antigens can not stimulate responses in neonatal mice or CBA/N mice that carry an X-linked immune B-cell defect (xid mice), whereas Type 1 antigens can.

Type 2 antigens induce weaker antibody responses as compared to T-dependent antigens such as proteins. Proteins are able to activate B-cells and induce the secretion of antibody by being processed into peptides and presented on the surface of the B-cell in the context of MHC class II, enabling the B-cell to interact with T-cells and receive additional signals required for maximal B-cell proliferation and maturation. However, whereas oligosaccharides may in some cases associate with MHC class II (Ishioka et al. J. Immunol. 148: 2446-2451) and lipidated

polysaccharides appear to associate with CD1 present on lymphocytes, (Fairhurst, R.M. et al. Immunology Today 19: 257 (1998)), there is no known mechanism of presentation for Type 2 antigens to T cells.

Nevertheless, the multiple repeat nature of the polysaccharide polymer
5 antigen can cause cross-linking of receptors on the B-cell surface, leading to B-cell activation by a mechanism not requiring T-cells. Thus polysaccharides are T-independent antigens and they are characterised in animals and human infants by the production of IgM antibodies, and the lack of boosting and immunological memory. It is only adult humans that can produce significant amounts of IgG antibody to
10 most (but not all) polysaccharide antigens. The ability to switch antibody isotype to IgG coincides with the appearance of the complement receptor 2 (CR2) on the B-cells of infants or toddlers between 1.5 to 2 years of age, and this may provide the additional signal required for activation and maturation of B-cells.

The present invention in one aspect provides a vaccine formulation which is
15 capable of raising an immune response to T independent antigen. Production of IgG antibodies to the capsular polysaccharides of bacteria is essential because the principal mechanism of protection against these bacteria, complement mediated lysis and opsonophagocytosis, are most effective with this antibody isotype (Maslanka et al. Clin Diag Lab Immunol 4: 156-67, and Romero-Steiner et
20 al. Clin Diag Lab Immunol 4: 415-22).

Polysaccharide antigen based vaccines are well known in the art, and four that have been licensed for human use include the Vi polysaccharide of *Salmonella typhi*, the PRP polysaccharide from *Haemophilus influenzae*, the tetravalent meningococcal vaccine composed of serotypes A, C, W135 and Y, and the 23-
25 Valent pneumococcal vaccine composed of the polysaccharides corresponding to serotypes 1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F, and 33. .

The latter three vaccines confer protection against bacteria causing respiratory infections resulting in severe morbidity and mortality in infants, yet
30 these vaccines have not been licensed for use in children less than two years of age because they are poorly immunogenic in this age group.

The licensed polysaccharide vaccines listed above have different demonstrated clinical efficacy. The Vi polysaccharide vaccine has been estimated to have an efficacy between 55% and 77% in preventing culture confirmed typhoid fever (Plotkin and Cam, Arch Intern Med 155: 2293-99). The meningococcal C polysaccharide vaccine was shown to have an efficacy of 79% under epidemic conditions (De Wals P, et al. Bull World Health Organ. 74: 407-411). The 23-valent pneumococcal vaccine has shown a wide variation in clinical efficacy, from 0% to 81% (Fedson et al. Arch Intern Med. 154: 2531-2535). The efficacy appears to be related to the risk group that is being immunised, such as the elderly, Hodgkin's disease, splenectomy, sickle cell disease and agammaglobulinemics (Fine et al Arch Intern Med. 154:2666-2677), and also to the disease manifestation. Pneumococcal pneumonia and Otitis media are diseases which do not have demonstrated protection by the 23-valent vaccine. It is generally accepted that the protective efficacy of the pneumococcal vaccine is more or less related to the concentration of antibody induced upon vaccination; indeed, the 23 polysaccharides were accepted for licensure solely upon the immunogenicity of each component polysaccharide (Ed. Williams et al. New York Academy of Sciences 1995 pp 241-249).

To increase the antibody response to the pneumococcal polysaccharides comprising the 23-valent vaccine, the present inventors tried to improve the immune response by addition of the immunostimulant QS21 EP 362 279 and dQS21 WO 96/33739; however, no increase in antibody responses to the polysaccharides in Rhesus monkeys could be measured.

Threadgill *et al* Vaccine 1998 Vol 16(1) p76 have recently reported that Immunostimulatory CpG oligonucleotides depress the polysaccharide specific antibody response when the oligonucleotide is formulated with *Pseudomonas aeruginosa* polysaccharide.

Surprisingly, the present inventors have found that it is possible to adjuvant the immune response to pneumococcal polysaccharide vaccines by formulating with an immunostimulatory CpG oligonucleotide, such formulations provided an immune response which produces significant levels of IgG antibodies.

According to the present invention there is provided a vaccine composition comprising a polysaccharide antigen adjuvanted by an Immunostimulatory oligonucleotide.

5 The polysaccharide antigen may be unconjugated or conjugated to a carrier protein such that it provides T-helper epitopes.

The oligonucleotides may be DNA or RNA, but preferably contain a hexamer motif: purine purine CpG pyrimidine pyrimidine. More preferably the internucleotide linkage are modified to increase stability of the oligonucleotide. Preferred modifications are phosphorothioate linkages. The *lytA* protein involved
10 in the catalytic degradation of the cell wall of pneumococci is produced at the time of autolysis, and is part of the competence induced operon (Mol. Microbiol 29:1125 (1998)). By definition the mRNA encoding *lytA* will be present in large quantities during synthesis of the *lytA* protein. Furthermore, the *lytA* protein contains a phosphoryl choline binding region that contains repeat DNA sequences (Yother and
15 Briles J Bacteriol. 174: 601 (1992)), and which may be found on many other choline binding proteins present in Streptococci. The following CpG sequences were identified from the phosphoryl choline binding regions of *lytA* and from choline binding protein A (*cbpA*) (Rosenow et al. Mol. Microbiol 25: 819-829 (1997)).

20

OLIGO 1: GCTACTGGTACG TACATTC AGACGGC TCTT (*lytA*)

OLIGO 2: ACTATCTAAACGCTAATGGTGCTATGGCGACAGGATGGCT (*cbpA*)

25 and may be utilised in the present invention.

The following oligonucleotide immunostimulatory sequences also form preferred embodiments of the invention.

OLIGO 3: TCC ATG ACG TTC CTG ACG TT

30 OLIGO 4: TCT CCC AGC GTG CGC CAT

The CpG and flanking sequences have been underlined, and there is conserved ACGT, ACG and GCG motifs. The sequences derived from the choline binding regions of pneumococcal proteins have two CpG motifs that repeated 10 or 15 nucleotide bases apart, and a motif based on this nucleotide base distance
5 between two CpGs occurs three times and five times respectively in the *lytA* and *CbpA* proteins. However, the published sequences have two CpG motifs that are seven or two nucleotide bases apart.

In one embodiment, when combined with commercially available 23 valent polysaccharide vaccine (Pneumovax, Pasteur Merieux), CpG adjuvantation
10 significantly augmented the immune response (IgG antibody) especially to polysaccharide types 19F and 14 when administered Intramuscularly.

Thus advantageously in an embodiment of the present invention it is possible to enhance the efficacy of a commercially available pneumococcal vaccine. This is particularly important in high risk populations, especially those which have sub-
15 optimal antibody responses to the polysaccharides. Such populations may include, but are not limited to, the elderly, patients with any of the following: splenectomy, congenital asplenia, hyposplenia, sickle cell disease, cyclic neutropenia, drug-induced neutropenia, aplastic anaemia, congenital agammaglobulinemia, hypogammaglobulinemia, selective IgG subclass deficiency, multiple myeloma,
20 chronic lymphocytic leukaemia, lymphoma, HIV infection, multifactorial conditions such as glucocorticoid treatment, malnutrition, cirrhosis of the liver, renal insufficiency, diabetes mellitus, alcoholism, chronic disease, hospitalisation, fatigue, stress, cold exposure, prior respiratory infection, influenza, asthma. It may also include healthy adults such as health workers, military trainees, prisoners, or
25 others including school attendees or travellers wishing to ensure full vaccine coverage.

In a preferred application, the CpG adjuvant is used to augment the response to the polysaccharide vaccine when used as booster in children between 6 and 24 months of age that have received their primary immunisation with a multivalent
30 pneumococcal polysaccharide-protein conjugate. Such vaccines utilised for primary immunisation may also advantageously, be adjuvanted with a CpG oligonucleotide. Accordingly in one embodiment there is provided a method of immunisation of a

patient comprising administering an effective amount of a vaccine according to the invention.

In a second embodiment there is provided a method of boosting an immune response to a subject previously primed to an antigen by administering a T-independent antigen with a CpG immunostimulatory oligonucleotide.

CpG adjuvantation may be applied, according to the present invention, to other polysaccharide and T-independent antigen based vaccines. These include, but are not limited to, the Vi polysaccharide vaccine against *Salmonella typhi*, the tetravalent meningococcal polysaccharide vaccine (comprising types A, C, W135 and Y), the polysaccharide and modified polysaccharides of group B meningococcus, polysaccharides from *Staphylococcus aureus*, polysaccharides from *Streptococcus agalactae*, polysaccharides from Mycobacteria, eg *Mycobacterium tuberculosis*, such as mannophosphoinositides trehaloses, mycolic acid, mannose capped arabinomannans, the capsule therefrom and arabinogalactans, polysaccharide from *Cryptococcus neoformans*, the lipopolysaccharides of non-typeable *Haemophilus influenzae*, the lipopolysaccharides of *Moraxella catharralis*, the lipopolysaccharides of *Shigella sonnei*, the lipopeptidophosphoglycan (LPPG) of *Trypanosoma cruzi*, the cancer associated gangliosides GD3, GD2, the tumor associated mucins, especially the T-F antigen, and the sialyl T-F antigen, and the HIV associated polysaccharide that is structurally related to the T-F antigen. Other T independent antigens may be derived from: Salmonella, Cholera, Echerichia, Chlamydia and T-independent antigens from Plasmodium.

Vaccine preparation is generally described in Pharmaceutical Biotechnology, Vol.6! Vaccine Design - the subunit and adjuvant approach, edited by Powell and Newman, Plenum Press, 1995. Encapsulation within liposomes is described, for example, by Fullerton, US Patent 4,235,877. Conjugation of proteins to macromolecules is disclosed, for example, by Likhite, US Patent 4,372,945 and by Armor et al, US Patent 4,474,757.

The amount of protein in each vaccine dose is selected as an amount which induces an immunoprotective response without significant, adverse side effects in typical vaccinees. Such amount will vary depending upon which specific immunogen is employed and how it is presented. Generally, it is expected that each

dose will comprise 0.1-1000 μ g of polysaccharide or polysaccharide - protein conjugate, preferably 2-100 μ g, most preferably 4-40 μ g. An optimal amount for a particular vaccine can be ascertained by standard studies involving observation of appropriate immune responses in subjects. Following an initial vaccination,
5 subjects may receive one or several booster immunisation adequately spaced.

The oligonucleotides utilised in the present invention are typically deoxynucleotides. In a preferred embodiment the internucleotide bond in the oligonucleotide is phosphorodithioate, or more preferably phosphorothioate bond, although phosphodiester are within the scope of the present invention. Other
10 internucleotide bonds which stabilise the oligonucleotide may be used.

The CpG oligonucleotides utilised in the present invention may be synthesized by any method known in the art (eg EP 468 520) conveniently such oligonucleotides can be synthesized utilising an automated synthesizer. Methods for producing phosphorothioate oligonucleotides or phosphorodithioate are
15 described in US 5,663,153, US 5,278,302 and WO 95/26204.

Example 1 - CpG adjuvantation of 23-valent Pneumococcal Polysaccharide in mice

20 Protection against pneumococcal infection is mediated by IgG antibody to the capsular polysaccharide, along with the deposition of complement that renders the bacteria susceptible to killing by neutrophils via opsonophagocytosis. Thus the protective efficacy of the vaccine may be estimated solely on the basis of IgG antibody induction. Groups of 10 mice were immunised once with a commercial
25 23-Valent pneumococcal polysaccharide vaccine at 1/10, 1/50 or 1/250 human dose (57.7, 11.5 and 2.3 μ g total polysaccharide respectively), and adjuvanted with CpG (50 μ g of oligo 1), CpG + Alum. Following immunization, serum IgG concentrations to the 4 most important serotype polysaccharides (6B, 14, 19F and 23F) were measured by ELISA every 7 days for 4 weeks.

Materials and Methods

The following groups were immunised. (10 balb/c mice per group):

23 Valent at 2.3, 11.5 and 57.5 µg/dose (1/250, 1/50 and 1/10 human dose)

5 23 Valent + CpG (50 µg) in the same dose range

23 Valent + CpG + Al(OH)₃ in the same dose range

Components used

Component	Batch	Concentration µg/ml	Buffer
23 Valent from Pasteur Mérieux (Pneumovax 23)	95K03-HC56630	1150	Saline
CpG	Oligo 3	5000	H ₂ O
Al(OH) ₃	96A0089	10380	H ₂ O

10

Formulation process

The Pneumovax was diluted in H₂O and 10-fold concentrated 10mM PO₄, 150 mM NaCl pH 6.8 to obtain 2.3, 11.5 or 57.7 µg of antigen per dose. CpG was added
 15 for 30min and for groups containing Al(OH)₃ the formulations were adsorbed for 30 min on either Al(OH)₃ (50µg). Thiomersal (50 µg/ml) was added as preservative.

ELISA

20 There were 10 animals per group, but since bleeds were performed every week, only 5 animals per week were bled. ELISA and opsonophagocytosis were performed on pooled sera.

25 The ELISA was performed to measure murine IgG using the protocol was derived from the WHO Workshop on the ELISA procedure for the quantitation of IgG antibody against *Streptococcus pneumoniae* capsular polysaccharides in human

serum. In essence, purified capsular polysaccharide is coated directly on the microtitre plate. Serum samples are pre-incubated with the cell-wall polysaccharide common to all pneumococcus and which is present in ca. 0.5% in pneumococcal polysaccharides purified according to disclosure (EP72513 B1). Jackson

5 ImmunoLaboratories Inc. reagents were employed to detect bound murine IgG. The titration curves were referenced to internal standards (monoclonal antibodies) modeled by logistic log equation. The calculations were performed using SoftMax Pro software. The maximum absolute error on these results expected to be within a factor of 2. The relative error is less than 30%.

10

Results

IgG isotype antibodies were found against serotypes 14 and 19G, but not against 6B and 23F, and the results for serotype 14 are presented in figure 1. The response
15 was dose dependent with 1/10 human dose giving the highest response, indicating that the IgG response was specific for the polysaccharide. This is unusual since mice normally only produce IgM against pneumococcal polysaccharides. The peak response was on day 14 post immunisation, which is not unusual since T-independent antigens do not induce memory.

20

Additional individual analysis were carried out to determine the variance and the statistical significance (data not shown). The response to 1/10 human dose 23-valent was (statistically) significantly increased when adjuvanted with CpG alone (for Type 19, GMC 0.8 compared to 3.7 $\mu\text{g/ml}$ $p=0.07$; for Type 14, GMC 0.19 compared
25 to 3.4 $\mu\text{g/ml}$, $p=.001$). This was also true for 1/50 and 1/250 doses when measured against type 14. In addition, responses were significantly increased to type 14 when adjuvanted with CpG + Alum.

The highest response was induced when the vaccine was adjuvanted with CpG
30 alone.

Example 2 - Effect of CpG adjuvantation on the immunogenicity of tetravalent Pneumococcal PS-PD conjugates in the infant rat model

The infant rat model was selected since published data showed that the relative immunogenicity of 4 pneumococcal polysaccharide protein conjugates in human infants was more similar to rats than mice. That is 6B < 23F < 14 < 19F for infant rats. Infants rats were selected because their immune system may have developmental immaturity similar that that found in human infants.

Infant rats were immunized with Clinical grade lots of Tetravalent pneumococcal polysaccharide-PD^(a) conjugates in a 5-fold dose range and with the adjuvants CpG and AlPO₄ + CpG. Oligo 1 was used at a dosage of 100µg. Animals were first immunized when they were 7 days old and received subsequent immunizations 14 and 28 days later. Serology was performed on samples from day 42 (14 days post III) and 56 (28 days post III).

The best adjuvant was CpG alone : it increased geometric mean IgG concentrations and opsonic titers to 6B, 23F and 19F, whereas titers for serotype 14 were comparable to the other adjuvanted preparations. The CpG alone formulation was also able to significantly increase the seroconversion rates to the 6B-PD serotype.

Materials and Methods

Vaccine groups

The vaccine lot DSP0401x contains Tetravalent PS-PD Clinical-grade lots D6BPJ208 + D14PDJ202 + D19PJ206 + D23PDJ212. ESPiL001 contains Tetravalent PS-LPD lots E6BL040P + E14L66P + E19FL033P + E23FL21P.

Group	Vaccine Lot	Adjuvant	Dose (µg each PS)
1	none	CpG	-
2	DSP0401x	none	0.1
3	DSP0401x	none	0.5
4	DSP0401x	AlPO ₄	0.1

Group	Vaccine Lot	Adjuvant	Dose (μ g each PS)
5	DSP0401x	AlPO ₄	0.5
6	DSP0401x	AlPO ₄	2.5
7	ESPL001	AlPO ₄	0.1
8	ESPL001	AlPO ₄	0.5
9	ESPL001	AlPO ₄	1.25
10	DSP0401x	CpG	0.1
11	DSP0401x	CpG	0.5
12	DSP0401x	CpG/AlPO ₄	0.1
13	DSP0401x	CpG/AlPO ₄	0.5

Components used

Component	Batch	Concentration μ g/ml	Buffer
Conjugate PD6B	D6BPDJ208	206	NaCl 0.2M pH 6.5
Conjugate PD14	D14PDJ202	186	NaCl 0.2M pH 6.5
Conjugate PD19	D19PDJ206	175	NaCl 0.2M pH 6.5
Conjugate PD23	D23PDJ212	158	NaCl 0.2M pH 6.5
monovalent PD6B	D6BPDD208	100	NaCl 150mM pH 6.1
monovalent PD14	D14PDD202	100	NaCl 150mM pH 6.1
monovalent PD19	D19PDD206	100	NaCl 150mM pH 6.1
monovalent PD23	D23PDD212	96	NaCl 150mM pH 6.1
monovalent LPD6B	E6BL040P	50	NaCl 150mM pH 6.1
monovalent LPD14	E14FL66P	50	NaCl 150mM pH 6.1
monovalent LPD19	E19FL033P	50	NaCl 150mM pH 6.1
monovalent LPD23	E23FL21P	50	NaCl 150mM pH 6.1
Tetavalent LPD	ESPL001	5/valence	NaCl 150mM pH 6.1
CpG	Oligo 1, WD1001	5000	H ₂ O
AlPO ₄	97D0045	5040	NaCl 150mM pH 6.1

Formulation Process**Non adsorbed tetravalents.**

- 5 The four conjugates are diluted in H₂O and 10-fold concentrated NaCl 150mM. Phenoxyethanol (500 µg/ml) is added as preservative.

If CpG is needed, the oligonucleotide is added to the non adsorbed tetravalent. The isotonicity and the dilution when needed are ensured by NaCl.

10

Adsorbed tetravalents

The four concentrated, adsorbed monovalents are diluted in H₂O and 10-fold concentrated 150mM NaCl before addition of the complement of AlPO₄.

- 15 Phenoxyethanol (500 µg/ml) is added as preservative.

If dilutions are needed, the tetravalents are diluted in AlPO₄ at 1 mg/ml. These diluents are prepared in NaCl 150 mM.

- 20 If CpG is needed, the oligonucleotide is added to the adsorbed tetravalent. The isotonicity is ensured by addition of NaCl 1500 mM and if dilutions are required, diluents of AlPO₄ at 1.3 or 1.8 mg/ml in NaCl are added.

All the formulations are prepared in non siliconized glass vials.

25

Immunisation Protocol

- Infant rats were randomised to different mothers and were 7 days old when they received the first immunisation. They received subsequent immunisations 14 and 28 days later. Bleeds were performed on day 42 (14 days post III) and 56 (28 days post III). All vaccines were injected s.c., and there were 10 rats per vaccine group.

ELISA

The ELISA was performed to measure rat IgG using the protocol derived from the WHO Workshop on the "ELISA procedure for the quantitation of IgG antibody against *Streptococcus pneumoniae* capsular polysaccharides in human serum". In essence, purified capsular polysaccharide is coated directly on the microtitre plate. Serum samples are pre-incubated with the cell-wall polysaccharide common to all pneumococcus and which is present in ca. 0.5% in pneumococcal polysaccharides purified. Jackson ImmunoLaboratories Inc. reagents were employed to detect bound rat IgG. The titration curves were referenced to the titration curve of a reference serum modeled by logistic log equation. The calculations were performed using SoftMax Pro software. The standard sera were calibrated using a method of corollary response, and the values were demonstrated to correspond to estimations of Ig concentrations found by immunoprecipitation (Ref. 21).

Opsonophagocytosis

The opsonophagocytic assay was performed following the CDC protocol (*Streptococcus pneumoniae* Opsonophagocytosis using Differentiated HL60 cells, version 1.1). Modification included the use of in-house pneumococcal strains, and the phagocytic HL60 cells were replaced by purified human PMN. Rat polyclonal sera were included as a positive control.

Results

Figure 2 shows the geometric mean IgG concentrations elicited against serotype 6B by the tetravalent combinations described in the materials and methods. For clarity, the axes are divided by adjuvant and dose. Similar results were obtained against the serotypes 19F and 23F, but type 14 had a more uniform response to all adjuvants and doses.

The biological activity of the pooled antisera from each adjuvant group and dose was measured by opsonophagocytosis. The opsonic activity relative to the concentration of IgG will give an estimate of the functional activity of the antisera.

- 5 The values, shown in Table 1 show that all adjuvants induce antibody that has approximately the same capacity to opsonise pneumococci. Thus CpG aids in the induction of specific antibody, and increases in antibody concentration correlate with increases in protective efficacy.

10 Conclusion

- AlPO4 (compared to no adjuvant) significantly increases the seroconversion rate, geometric mean IgG concentration, opsonic activity and immunological
15 memory to tetravalent PS-PD.
- The 0.1 µg dose is significantly more immunogenic than 0.5 µg dose for serotypes 6B, 19F and 23F PS-PD conjugates on AlPO4.
20
- IgG concentrations are significantly increased against serotypes 6B, 19F and 23F when the conjugate vaccine is adjuvanted with CpG compared to AlPO4. This is confirmed by increased seroconversion rates and increased
25 opsonophagocytic titres.

Table 1. Relative opsonic activity (Concentration of IgG required for 50% killing of pneumococcus) compared by serotype and adjuvant.

Vaccine	Adjuvant	Dose	Concentration of IgG required for 50% killing				Average by adjuvant
		μg	6B	14	19F	23F	
DSP0401x	none	0,1	0,32	0,30	0,30	0,37	0.26±0.14
		0,5	No Value	0,015	No Value	No Value	
DSP0401x	AIPO4	0,1	0,02	0,31	0,40	0,09	0.20±0.15
		0,5	No Value	0,05	0,22	No Value	
		2,5	No Value	0,32	#VAL	No Value	
ESPL001	AIPO4	0,1	0,08	0,46	No Value	0,22	0.35±0.27
		0,5	0,11	0,71	0,75	0,08	
		1,25	0,10	0,55	0,66	0,20	
DSP0401x	CPG	0,1	0,42	0,15	No Value	0,20	0.24±0.10
		0,5	0,21	0,30	No Value	0,17	
DSP0401x	CPG / AIPO4	0,1	0,27	0,10	No Value	0,21	0.20±0.14
		0,5	No Value	0,10	0,44	0,09	
Average	by serotype		0.19±0.14	0.29±0.20	0.45±0.20	0.18±0.09	

5

Example 3 - Effect of CpG adjuvantation on the immunogenicity of 11-Valent Pneumococcal PS-PD conjugates in the infant rat model

- 10 Example 2 showed that CpG adjuvantation of conjugate vaccines resulted in fold increases of the order of 5 to 10 times higher than with conventional adjuvants (Aluminium). In order to determine whether these effects were dependent on the Oligo sequence, dosage, or formulation, further experimentation was undertaken.
- 15 CpG OLIGO 2 was selected and used at a lower dosage, that is 1 and 10 μg . It was also adsorbed onto $\text{Al}(\text{OH})_3$, and combined with the conjugate vaccines.

In addition, since the immunological characteristics of each polysaccharide may be different, 11 serotypes were tested.

Material and Methods

Table 2. Choice of pneumococcal PS-PD lots

Serotype	1	3	4	5	6B	7F	9V	14	18C	19F	23F
Lot number	017	040	218	024	209	019	222	204	221	207	213

Formulation

To examine the effect of different advanced adjuvants, the dosage of conjugate was held constant at 0.1 µg of each polysaccharide, and the adjuvants AlPO₄, Al(OH)₃ and CpG were formulated in different dosages and combinations. In total, 10 different combinations were tested, including no adjuvant at all. These are listed numerically in Table 3 for reference.

Preparation of diluents

Two diluents were prepared in NaCl 150 mM/phenoxy

A: AlPO₄ at 1mg/ml.

B: CpG on Al(OH)₃ at 200 and 1000 µg/ml respectively. Weight ratio CpG/Al(OH)₃ = 1/5

Preparation of adsorbed undecavalent

The eleven concentrated, adsorbed PS-PD monovalents were mixed at the right ratio. The complement of AlPO₄ was added. When needed, CpG (CpG adsorbed on Al(OH)₃) or diluent was added.

Preparation of non-adsorbed undecavalent

The eleven PS-PD conjugates were mixed and diluted at the right ratio in NaCl 150 mM pH 6.1, phenoxo. When needed, CpG was added either as a solution (non adsorbed) or as CpG adsorbed on Al(OH)₃.

The formulations for all injections were prepared 18 days before the first administration.

5 **Table 3. Summary Table of Adjuvant Formulations tested with 11-Valent Pneumococcal PS-PD in Infant Rats**

Group	AlPO4	CpG	Al(OH)3	Description
1				None
2	100			AlPO4
3		1		CpG low
4		10		CpG High
5		1	4.5	CpG ads low
6		10	50	CpG ads high
7	100	1		CpG low Conj ads
8	100	10		CpG Hi Conj ads
9	95	1	4.5	CpG&Conj ads low
10	50	10	50	CpG&Conj ads Hi

Immunisation Protocol

10

Infant OFA rats were randomised to different mothers and were 7 days old when they received the first immunisation. They received 2 additional immunisations 14 and 28 days later. A bleed as performed on day 56 (28 days post III). All vaccines were injected s.c., and there were 10 rats per vaccine group.

15

ELISA

The ELISA was performed to as described in example 2.

20 Opsonophagocytosis

The opsonophagocytic assay was performed following the CDC protocol (Streptococcus pneumoniae Opsonophagocytosis using Differentiated HL60 cells, version 1.1). Modification included the use of in-house pneumococcal strains, and
 25 the phagocytic HL60 cells were replaced by purified human PMN. In addition, 3 mm glass beads were added to the microtitre wells to increase mixing, and this allowed reduction of the phagocyte:bacteria ratio which was recommended to be 400.

Results

Tables 4 to 7 below show the geometric mean IgG concentration, seroconversion rate and arithmetic mean opsonophagocytic titre determined for 4 serotypes of pneumococci after immunisation with an 11Valent pneumococcal PS-Protein D conjugate vaccine adjuvanted with different formulations of CpG OLIGO 2. Compared to no adjuvant, 10 µg CpG induced significant higher IgG concentrations for all serotypes. CpG induced significantly higher IgG concentrations than AlPO₄ for serotypes 1, 6B, 18C and 19F.

10

For comparison, included in the Tables are the results from Example2 using OLIGO 1. There are no significant differences in the IgG responses induced by the two OLIGO sequences when OLIGO 2 is used at 10 µg. However, OLIGO 2 at 1 µg shows no immunostimulatory effects evidenced in that the induced IgG concentrations are not significantly different from without CpG.

15

Adsorption of OLIGO 2 on Al(OH)₃ reduces the immunostimulatory effect, and the induction of antibody is not significantly different than AlPO₄ as adjuvant.

TABLE 4

Serotype 6B Geometric Mean IgG Concentration, Seroconversion, and Mean Opsonic Titre on Day 28 Post III Immunisation of Infant Rats with 11-Valent PS-PD using Different Adjuvants (And Comparison with Tetravalent Immunisation, Example 2)

Gro up	Al PO 4 µg	Oligo 1 µg	Oligo 2 µg	6B GMC IgG µg/ml	6B Sero- con- versio n	6B Opso Titre*	6B GMC IgG µg/ml	6B Sero- con- version	6B Opso Titre*
				Example 2			Example 3		
1				0.047	2/10	12.5	0.004	1/10	<6.25
2	100			0.048	4/10	65	0.019	4/10	<6.25
3			1				0.003	1/10	<6.25
4			10				1.682	10/10	157
		100		0.63	8/10	48			
5			1 µg on Al(OH) ₃				0.015	6/10	<6.25
6			10 µg on Al(OH) ₃				0.007	3/10	<6.25
7	100		1				0.029	7/10	<6.25
8	100		10				0.469	9/10	77
	100	100		0.46	7/10	75			
9	95		1 µg on Al(OH) ₃				0.040	5/10	38
10	50		10 µg on Al(OH) ₃				0.022	7/10	<6.25

TABLE 5

5 Serotype 14 Geometric Mean IgG Concentration, Seroconversion, and Mean Opsonic Titre on Day 28 Post III Immunisation of Infant Rats with 11-Valent PS-PD using Different Adjuvants (And Comparison with Tetravalent Immunisation, Example 2)

Gro up	Al PO 4 µg	Oligo 1 µg	Oligo 2 µg	14 GMC IgG µg/ml	14 Sero- con- versio n	14 Opso Titre*	14 GMC IgG µg/ml	14 Sero- con- version	14 Opso Titre*
				Example 2			Example 3		
1				0.046	3/10	64	0.022	3/10	<6.25
2	100			0.99	10/10	88	0.237	8/10	27
3			1				0.035	4/10	<6.25
4			10				0.361	10/10	88
		100		0.66	9/10	295			
5			1 µg on Al(OH) ₃				0.093	9/10	<6.25
6			10 µg on Al(OH) ₃				0.155	9/10	27
7	100		1				0.134	7/10	<6.25
8	100		10				2.028	10/10	188
	100	100		2.3	10/10	888			
9	95		1 µg on Al(OH) ₃				0.140	6/10	138
10	50		10 µg on Al(OH) ₃				0.196	10/10	<6.25

TABLE 6

5 Serotype 19F Geometric Mean IgG Concentration, Seroconversion, and Mean Opsonic Titre on Day 28 Post III Immunisation of Infant Rats with 11-Valent PS-PD using Different Adjuvants (And Comparison with Tetravalent Immunisation, Example 2)

Gro up	Al PO 4 µg	Oligo 1 µg	Oligo 2 µg	19F GMC IgG µg/ml	19F Sero- con- versio n	19F Opso Titre*	19F GMC IgG µg/ml	19F Sero- con- version	19F Opso Titre*
				Example 2			Example 3		
1				0.04	2/10	64	0.021	2/10	<6.25
2	100			1.07	9/10	367	0.222	7/10	79
3			1				0.015	3/10	<6.25
4			10				4.287	10/10	415
		100		12.	10/10	>1600			
5			1 µg on Al(OH) ₃				0.417	9/10	32
6			10 µg on Al(OH) ₃				1.612	9/10	94
7	100		1				0.441	10/10	135
8	100		10				9.475	10/10	>1600
	100	100		11.0	10/10	>1600			
9	95		1 µg on Al(OH) ₃				0.438	9/10	377
10	50		10 µg on Al(OH) ₃				0.258	7/10	165

TABLE 7

5 Serotype 23F Geometric Mean IgG Concentration, Seroconversion, and Mean Opsonic Titre on Day 28 Post III Immunisation of Infant Rats with 11-Valent PS-PD using Different Adjuvants (And Comparison with Tetravalent Immunisation, Example 2)

Gro up	Al PO 4 µg	Oligo 1 µg	Oligo 2 µg	23F GMC IgG µg/ml	23F Sero- con- versio n	23F Opso Titre*	23F GMC IgG µg/ml	23F Sero- con- version	23F Opso Titre*
				Example 2			Example 3		
1				0.06	2/10	<6.25	0.152	3/10	<6.25
2	100			0.29	10/10	70	0.56	8/10	<6.25
3			1				0.114	4/10	<6.25
4			10				1.305	9/10	192
		100		2.0	10/10	454			
5			1 µg on Al(OH) ₃				0.28	7/10	<6.25
6			10 µg on Al(OH) ₃				0.107	2/10	<6.25
7	100		1				0.243	4/10	<6.25
8	100		10				1.545	9/10	862
	100	100		1.1	10/10	265			
9	95		1 µg on Al(OH) ₃				0.255	3/10	44
10	50		10 µg on Al(OH) ₃				0.331	6/10	<6.25

10

Example 5 Influence of CpG on Boosting with Polysaccharide after Priming with Polysaccharide-Conjugate Vaccines, and on Priming with Polysaccharide.

15 The previous examples have demonstrated the ability of CpG to adjuvant the immune response to T-independent antigens, and to T-independent antigens coupled to a protein carrier. There remained to be considered whether CpG could adjuvant a memory response elicited by boosting with a T-independent antigen after priming with T-dependent antigen. It was of further interest to determine if CpG could act

20 to induce priming by a T-independent antigen.

To determine these effects, mice were primed with either pneumococcal polysaccharide, or CpG adjuvanted pneumococcal polysaccharide, or Protein D conjugate pneumococcal polysaccharide.

5

Immunisation Protocol

Six to 8 week old balb/c mice were immunise subcutaneously with the vaccine formulations described below. The dosage was 1 μ g per polysaccharide for both conjugated and non-conjugated formulations. A test bleed was performed 14 days later to measure IgG concentrations. After 56 days, another test bleed was performed, and then a booster vaccination was given, and a final test bleed was performed 14 days later, that is 70 days after the first immunisation.

Group	Prime	Boost
1	Saline	Conjugate
2	PS	PS
3	PS/Cp	PS
4	PS	Conjugate
5	PS/Cp	Conjugate
6	Conjug	PS
7	Conjug	PS/CpG
8	Conjug	Conjugate

15

Components used

Component	Batch	Conc μ g/ml	Buffer	Adsorption	PS conc after adsorption μ g/ml	Buffer after adsorption
PS6b	6b/24	2000	NaCl 150mM			
PS14	14/19	2000	NaCl 150mM			
PS19	19f/26b	2000	NaCl 150mM			
PS23	23f/29	2000	NaCl 150mM			
Conjugate PDPS6B	D6BPDJ 209			MBSP9801	100	NaCl 150mM pH 6.1/phenoxo

Component	Batch	Conc $\mu\text{g/ml}$	Buffer	Adsorption	PS conc after adsorption $\mu\text{g/ml}$	Buffer after adsorption
Conjugate PDPS14	D14PDJ 204			MBSP9801	100	NaCl 150mM pH 6.1/phenoxy
Conjugate PDPS19	D19PDJ 207			MBSP9801	100	NaCl 150mM pH 6.1/phenoxy
Conjugate PDPS23	D23PDJ 213			MBSP9801	100	NaCl 150mM pH 6.1/phenoxy
CpG	Oligo 1	5000	H ₂ O			
St Pn AlPO ₄ diluent	97D004 5				1000	NaCl 150mM pH 6.1/phenoxy

Formulation process

Preparation of 4 concentrated, adsorbed monovalents (PS-PD conjugates)

5

The concentrated, adsorbed monovalents were prepared according the procedure described above in Example 2.

Preparation of tetravalent (PS-PD conjugates)

10

The four concentrated, adsorbed monovalents were mixed at the right ratio (1 μg of each valence/dose) and diluted in NaCl pH6.1. The complement of AlPO₄ (10 μg /dose) was added as a diluent at 1mg/ml in NaCl 150mM pH6.1 containing 5mg/ml of phenoxyethanol.

15

Preparation of non-conjugated, non-adsorbed, tetravalent with or without CpG (free PS)

The four free PS were mixed at the right ratio (1 μg of each valence/dose) and diluted in NaCl pH6.1. When needed, CpG (100 μg /dose) was added. Five mg/ml of phenoxyethanol were added as preservative.

20

The formulations for both injections were prepared 6 days before the first administration in non siliconized glass vials.

Formulation process

5

Preparation of 4 concentrated, adsorbed monovalents (PS-PD conjugates)

The concentrated, adsorbed monovalents were prepared according the procedure describe above.

10 **Preparation of tetravalent (PS-PD conjugates)**

The four concentrated, adsorbed monovalents were mixed at the right ratio (1 μ g of each valence/dose). The complement of AlPO₄ (10 μ g/dose) was added as a diluent at 1mg/ml in NaCl 150mM pH6.1 containing 5mg/ml of phenoxyethanol.

15

Preparation of non conjugated, non adsorbed, tetravalent with or without CpG (free PS)

20 The four free PS were mixed at the right ratio (1 μ g of each valence/dose) and diluted in NaCl pH6.1. When needed, CpG was added. Five mg/ml of phenoxyethanol were added as preservative.

The formulations for both injections were prepared 6 days before the first administration in non siliconized glass vials.

25

ELISA

The ELISA was performed as described in Example 1

Results

The results of this experiment are the priming and the boosting. The results of the priming were consistent with previous observations (Example 1) in that increased seroconversion and higher IgG concentrations were found in mice that were immunised with CpG adjuvanted polysaccharide compared to plain polysaccharide. As was found in Example 1, the increases in type 14 IgG concentration with CpG adjuvantation are statistically significant compared to PS alone, and the increases for type 19F approach significance. However, the IgG concentrations with CpG adjuvantation were not as high as observed in Example 1. To explain this difference, only two differences in the experiments were made, the valency of the vaccine (23 valent versus 4 valent) and the route of immunisation (intramuscular versus subcutaneous). Since the reduction of valence is not expected to decrease immunogenicity, the evidence indicates that the route of immunisation is important for optimal CpG adjuvantation of T-independent antigens. This is consistent with a recent publication which disclosed a failed attempt to use CpG adjuvantation of a plain polysaccharide vaccine. The route of immunisation employed was interperitoneal (Threadgill et al Vaccine 1998 Vol 16(1) p76).

	Seroconversion	GMC
PS 14	2/20*	0.07
PS14/CpG	12/20* δ	0.15
Conjugate	24/30 δ	1.04
PS19F	1/20 ξ	0.08
PS19F CpG	4/20 $\xi\varpi$	0.10
Conjugate	22/30 ϖ	0.35

* p = 0.001 Fisher's exact test

δ p = 0.11 Fisher's exact test

ξ p = 0.17 Fisher's exact test

ϖ p < 0.001 Fisher's exact test

- In the second part of this experiment, animals primed with either PS, PS/CpG or conjugate vaccine, were boosted with PS, or with PS/CpG or with conjugate. To normalise the data for comparison, the fold increase in IgG was determined 14 days after the booster was given, and the number of animal showing an increase in antibody concentration were counted as responders.

Prime	Boost	Geometric Fold Increase	Positive responders
PS	PS	1.7*	5/10
PS/CpG	PS	2.8*	6/10
Conjugate	PS	0.78 ξ	1/10 δ
Conjugate	PS/CpG	1.7 ξ	6/10 δ
Conjugate	Conjugate	4.2	7/10

- * p = 0.09 Student's t-test
 ξ p = 0.12 Student's t-test
 δ p = 0.03 Fisher's exact test

Discussion

- This example confirms the results presented in Example 1, but has revealed that the mode of immunisation may be important for optimal immunity. In an extension of the experiment to boosting and memory, two interesting characteristics of CpG adjuvantation are demonstrated. The first is that priming with PS adjuvanted with CpG leads to a higher fold increase upon boosting with polysaccharide, and there is a trend towards statistical significance. This would indicate that CpG was able to induce better memory. The second characteristic is that CpG can adjuvant a memory response induced by polysaccharide in animals primed with conjugate vaccine.

Conclusions

CpG is able to induce in mice an antibody isotype switch against non-conjugated
5 polysaccharides. The magnitude of the IgG response is higher with CpG.

Claims

1. A formulation comprising a CpG oligonucleotide and T-independent type 1 or type 2 antigens or polysaccharide conjugate antigen.
- 5 2. A formulation as claimed in claim 1, wherein the antigen is selected from the unconjugated Streptococcus Pneumoniae Polysaccharide, unconjugated meningococcal Polysaccharide, unconjugated Salmonella polysaccharide, unconjugated group A or group B streptococcal polysaccharides, polysaccharides
10 from mycobacteria unconjugated Pseudomonas mucoid polysaccharide or TF-antigens derived from HIV, or lipopolysaccharides or detoxified lipopolysaccharides derived from Salmonella, Cholera, Echerichia, Neisseria, Chlamydia, Shigella, Pertussis, Haemophilus, or Pseudomonas, or T-independent antigens from Plasmodium.
- 15 3. A formulation as claimed in claim 1 & 2 wherein the CpG deoxyribo- or ribo-oligonucleotide has an internucleotide bond, selected from phosphodiester, phosphorodithioate and phosphorothioate.
- 20 4. A formulation comprising a CpG deoxyribo or ribo-oligonucleotide sequence claimed in claim 2 which contains two CpG sequences that are separated by seven or more nucleotide base pairs.
5. A deoxyribo or ribo-oligonucleotide sequence claimed in claim 4 which
25 contains two CpG sequences that are separated by 10 to 15 nucleotide base pairs.
6. A formulation as claimed herein wherein the CpG oligonucleotide is selected from the group:

30 GCTACTGGTACG TACATTC AGACGGC TCTT
ACTATCTAAACGCTAATGGTGCTATGGCGACAGGATGGCT

TCC ATG ACG TTC CTG ACG TT

TCT CCC AGC GTG CGC CAT

7. A vaccine composition as claimed herein for use in medicine.

5

8. A method of inducing an immune response to T independent type 1 or type 2 antigen or a polysaccharide conjugate antigen, said method comprising administering a safe and effective amount of a formulation as claimed herein to a patient.





Fig.1 anti-PS14 IgG levels (days 7-28) in sera B45124a

2/2

**Fig.2** B45124a

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 39/39, 39/02, 39/015 // A61K 39/09	A3	(11) International Publication Number: WO 99/33488 (43) International Publication Date: 8 July 1999 (08.07.99)
(21) International Application Number: PCT/EP98/08562 (22) International Filing Date: 18 December 1998 (18.12.98) (30) Priority Data: 9727262.9 24 December 1997 (24.12.97) GB (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM BIOLOGICALS S.A. [BE/BE]; Rue de l'Institut 89, B-1330 Rixensart (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): DALEMANS, Wilfried, L., J. [BE/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE). LAFERRIERE, Craig, Antony, Joseph [CA/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE). PRIEELS, Jean-Paul [BE/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE). (74) Agent: DALTON, Marcus, Jonathan, William; SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 2 September 1999 (02.09.99)
(54) Title: ADJUVANTED VACCINE FORMULATION (57) Abstract <p>The present invention provides vaccine formulations comprising T-independent or polysaccharide conjugate vaccines adjuvanted with an immunostimulatory CpG oligonucleotide.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/08562

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K39/39 A61K39/02 A61K39/015 //A61K39/09

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>R.S. CHU ET AL.: "CpG OLIGODEOXYNUCLEOTIDES ACT AS ADJUVANTS THAT SWITCH ON T HELPER 1 (Th1) IMMUNITY." JOURNAL OF EXPERIMENTAL MEDICINE, vol. 186, no. 10, 17 November 1997, pages 1623-1631, XP002910130 NEW YORK, N.Y., US see page 1624, right-hand column, line 1; figure 1; table 1 see page 1629, right-hand column, paragraph 4 - page 1630, right-hand column, paragraph 1</p> <p style="text-align: center;">--- -/-</p>	1-8

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 June 1999

Date of mailing of the international search report

06/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Ryckebosch, A

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 98/08562

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	A.M. KRIEG ET AL.: "CpG MOTIFS IN BACTERIAL DNA TRIGGER DIRECT B-CELL ACTIVATION." NATURE, vol. 374, 6 April 1995, pages 546-549, XP002106695 LONDON, GB cited in the application see page 548, right-hand column, paragraph 2 see page 549, left-hand column, paragraph 2	1-8
P,X	--- D.S. THREADGILL ET AL.: "MITOGENIC SYNTHETIC POLYNUCLEOTIDES SUPPRESS THE ANTIBODY RESPONSE TO A BACTERIAL POLYSACCHARIDE." VACCINE, vol. 16, no. 1, January 1998, pages 76-82, XP002911830 GUILDFORD, GB cited in the application see the whole document	1-3,7,8
P,X	--- EP 0 855 184 A (G.B. LIPFORD ET AL.) 29 July 1998 see page 4, line 21 - page 5, line 31; claims; table 4	1,3,7,8
P,A	--- WO 98 37919 A (UNIVERSITY OF IOWA RESEARCH FOUNDATION) 3 September 1998 see SEQ ID NO:27 see claims; table 4 -----	1-8

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 98/08562

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim 8 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims: it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/08562

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 855184 A	29-07-1998	AU 6293498 A	18-08-1998
		WO 9832462 A	30-07-1998
WO 9837919 A	03-09-1998	AU 6667498 A	18-09-1998